

The Geoengineering Model Intercomparison Project (GeoMIP)

Ben Kravitz,^{1*} Alan Robock,¹ Olivier Boucher,^{2†} Hauke Schmidt,³ Karl E. Taylor,⁴ Georgiy Stenchikov^{1,5} and Michael Schulz⁶

¹Department of Environmental Sciences, Rutgers University, New Brunswick, NJ, USA

²Met Office Hadley Centre, Exeter, UK

³Max Planck Institute for Meteorology, Hamburg, Germany

⁴Program for Climate Model Diagnosis and Intercomparison, Lawrence Livermore National Laboratory, Livermore, CA, USA

⁵King Abdullah University of Science and Technology, Thuwal, Saudi Arabia

⁶Laboratoire des Sciences du Climat et de l'Environnement, Gif-sur-Yvette, France

*Correspondence to:

Ben Kravitz, Rutgers University,
Environmental Sciences, 14
College Farm Road, New
Brunswick, NJ 08901, USA.

E-mail:

benkravitz@envsci.rutgers.edu

†The contribution of O. Boucher was written in the course of his employment at the Met Office, UK, and is published with the permission of the Controller of HMSO and the Queen's Printer for Scotland.

Abstract

To evaluate the effects of stratospheric geoengineering with sulphate aerosols, we propose standard forcing scenarios to be applied to multiple climate models to compare their results and determine the robustness of their responses. Thus far, different modeling groups have used different forcing scenarios for both global warming and geoengineering, complicating the comparison of results. We recommend four experiments to explore the extent to which geoengineering might offset climate change projected in some of the Climate Model Intercomparison Project 5 experiments. These experiments focus on stratospheric aerosols, but future experiments under this framework may focus on different means of geoengineering. Copyright © 2011 Royal Meteorological Society and Crown Copyright

Keywords: geoengineering; climate modeling; CMIP5; model evaluation; SRM; monsoon

Received: 26 February 2010
Revised: 28 September 2010
Accepted: 8 November 2010

1. Introduction

Since the idea of geoengineering was thrust back into the scientific arena by Crutzen (2006) and Wigley (2006), many have wondered whether it could reduce global warming as mitigation measures are implemented. Several methods of geoengineering have been discussed, but Lenton and Vaughan (2009) argue that among the most feasible is through stratospheric sulphate aerosols. Analyses by Robock *et al.* (2009) indicate that such a scenario would be relatively inexpensive, especially in comparison with the cost of mitigation as determined by the Intergovernmental Panel on Climate Change (IPCC, 2007b), potentially making this idea attractive to policy makers.

However, Robock *et al.* (2009) point out that stratospheric geoengineering with sulphate aerosols could have unintended and possibly harmful consequences, including potential impacts on the hydrologic cycle and ozone depletion. For policymakers to be able to make informed decisions, the strength and patterns of these climate system responses need to be understood, and climate modeling will play an important part in this analysis. So far, several groups have conducted experiments, but these largely cannot be

directly compared. For instance, Robock *et al.* (2008) and Rasch *et al.* (2008a) used a 5 Tg SO₂ per year injection rate into the tropical lower stratosphere, while Jones *et al.* (2010) injected the same amount, but uniformly globally. In contrast, Govindasamy and Caldeira (2000), Govindasamy *et al.* (2002, 2003), Matthews and Caldeira (2007), and Bala *et al.* (2008) reduced the solar constant to approximate the net effects of stratospheric aerosols on the planetary energy balance. Robock *et al.* (2008) and Jones *et al.* (2010) ramped up the anthropogenic greenhouse gas forcing using the IPCC A1B scenario (IPCC, 2007a), while the others conducted equilibrium simulations at 2×CO₂.

Bala *et al.* (2008) explained why globally averaged precipitation would be reduced if the solar constant is reduced to balance the radiative forcing from increased greenhouse gas concentrations. However, simulation of the spatial patterns of such a reduction would likely be model-dependent. The results of Robock *et al.* (2008) indicate that stratospheric geoengineering in order to compensate for increased greenhouse gas concentrations would reduce summer monsoon rainfall in Asia and Africa, potentially threatening the food supply for billions of people. Jones *et al.* (2010)

Table I. A summary of the four experiments included in this proposal.

G1	Instantaneously quadruple the CO ₂ concentration (as measured from pre-industrial levels) while simultaneously reducing the solar constant to counteract this forcing (Figure 1).
G2	In combination with a 1% increase in CO ₂ concentration per year, gradually reduce the solar constant to balance the changing radiative forcing (Figure 2).
G3	In combination with RCP4.5 forcing, starting in 2020, gradual ramp-up the amount of SO ₂ or sulphate aerosol injected, with the purpose of keeping global average temperature nearly constant (Figure 3). Injection will be done at one point on the Equator or uniformly globally. The actual amount of injection per year can be based on Hansen <i>et al.</i> (2005) but may need to be fine tuned to each model.
G4	In combination with RCP4.5 forcing, starting in 2020, daily injections of a constant amount of SO ₂ at a rate of 5 Tg SO ₂ per year at one point on the Equator through the lower stratosphere (~16–25 km in altitude) or the particular model's equivalent. These injections would continue at the same rate through the lifetime of the simulation (Figure 4).

got similar results, but Rasch *et al.* (2008a) found different regional patterns. Past large volcanic eruptions have disrupted the summer monsoon (Oman *et al.*, 2005; Trenberth and Dai, 2007) and even produced famine (Oman *et al.*, 2006), but direct comparisons between geoengineering with stratospheric sulphate aerosols and large volcanic eruptions are limited by the differences in forcing. Some unanswered questions include whether a continuous stratospheric aerosol cloud would have the same effect as a transient one and to what extent regional changes in precipitation would be compensated by regional changes in evapotranspiration. A consensus has yet to be reached on these, as well as other, important issues.

To answer these questions, we propose a suite of standardized climate modeling experiments to be performed by interested modeling groups. We also propose to establish a coordinating framework for performing such experiments, which will be known as the Geoengineering Model Intercomparison Project (GeoMIP). Aside from coordinating the experiments described here, GeoMIP may consider additional geoengineering experiments in response to interest from climate modeling groups and the broader community. The particular experiment suite outlined in this document consists of four experiments, all of which are relevant to the geoengineering strategy of injecting stratospheric sulphate aerosols in an attempt to offset greenhouse gas warming. The Program for Climate Model Diagnosis and Intercomparison (PCMDI) has consented to archive results from these experiments, so they can be openly studied. We anticipate that this set of standardized experiments will permit the level of intercomparison necessary to achieve confidence in the results, similar to the level of scientific consensus that is published in the assessment reports of the IPCC. Initially, largely for practical reasons, the number of simulations to be performed must be kept small because the Climate Model Intercomparison Project 5 (CMIP5; Taylor *et al.*, 2008) is already stretching the capabilities of the modeling groups.

2. Experiment design

We use the codes G1, G2, G3, and G4 to refer to the four simulations that will be conducted in this suite

of experiments. We summarize these four experiments in Table I. G1, G2, and G3 are designed to produce an annual mean global radiative balance at the top of the atmosphere. We seek to determine commonalities and differences among climate model responses to these particular schemes of geoengineering. G1 and G2 are the simplest possible explorations of balancing increased longwave forcing with reduced shortwave forcing, i.e. through a reduction of the solar constant. These idealized experiments are expected to reveal the basic model responses to this forcing balance without the added complication of differing treatments of stratospheric aerosols in the various models. The idealized specification of forcing also makes it especially easy to implement. In all of these experiments, we define radiative forcing to be the 'adjusted forcing', which applies after so-called 'fast' radiative responses (e.g. stratospheric adjustment) occur, as discussed, for example, in Hansen *et al.* (2005). We note that we will unlikely be able to attain a perfect balance in radiative forcings, but we are aiming for a net balance as close to zero as possible. Included in CMIP5 is a historical run, which will include simulation of past volcanic eruptions. We will use the results of these simulations to validate the models as a part of our interpretation of the GeoMIP runs.

The G1 experiment will be initiated from a model control run and will build on a CMIP5 simulation in which the CO₂ concentration is instantaneously quadrupled. We choose this experiment to ensure a high signal-to-noise ratio of the climate response to radiative forcing from CO₂. In G1, the global average radiative forcing from the CO₂ will be balanced by a reduction of the solar constant. The CO₂ radiative forcing will be measured during the CMIP5 quadrupled CO₂ run, and the reduction in solar constant needed to compensate for this forcing will be based on a simple calculation using global average planetary albedo. A correction to this first estimate of solar constant change can be made after simulating a few years and monitoring the radiative balance. If a correction is necessary, the simulation will be restarted from the control run. In each model, a different solar constant change may be needed as both the CO₂ radiative forcing and the planetary albedo may differ from one model to the next. The tuning procedure described above also determines the change in solar constant that will be applied in the




Figure 1. Schematic of experiment G1. The experiment is started from a control run. The instantaneous quadrupling of CO₂ concentration from pre-industrial levels is balanced by a reduction in the solar constant until year 50.




Figure 2. Schematic of experiment G2. The experiment is started from a control run. The positive radiative forcing of an increase in CO₂ concentration of 1% per year is balanced by a decrease in the solar constant until year 50.

G2 experiment. Figure 1 illustrates the net radiative balance that would result from G1.

Similar to G1, the G2 experiment will involve a reduction in solar forcing to counteract the additional forcing due to increasing CO₂ concentration. However, the G2 experiment will build on the CMIP5 run specifying a 1% per year increase in CO₂, starting from a model control run. In G2, the global average radiative forcing from increases in CO₂ concentration will be balanced by gradually reducing the solar constant. As we prescribe an exponentially increasing CO₂ concentration, and the radiative forcing scales with the logarithm of CO₂ concentration, the solar constant will be prescribed to decrease linearly over time, with the scaling for the solar constant changes inferred from G1. Figure 2 illustrates the radiative balance that would result from G2.

Experiment G3 is similar to G1 and G2 but more realistic, in the sense that it will provide a scenario of possible implementation of stratospheric geoengineering (Figure 3). It assumes an RCP4.5 scenario (representative concentration pathway, with a radiative forcing of 4.5 W m⁻² in the year 2100; Moss *et al.*, 2008), but with additional stratospheric aerosol added starting in the year 2020, which is a reasonable estimate of when the delivery systems needed to inject the aerosols might be ready. Stratospheric aerosols will be added gradually, balancing the anthropogenic forcing to keep the planetary temperature nearly constant. The aim of this experiment is to achieve an ongoing radiative balance, which will likely require differing amounts of aerosol, with a time-varying size distribution, to be added in the various models. Ideally, the models will create, grow, and transport sulphate aerosols from an equatorial injection of SO₂. If a model does not have this capability, aerosols can be added at the Equator or globally in a way similar to each model's treatment of volcanic aerosols. If the model is capable, inclusion of O₃ chemistry or the carbon cycle, as well as the relevant couplings with

the physical climate system, will allow additional scientific issues to be addressed, but the models should be run in concentration-driven rather than emission-driven mode for the carbon cycle. The G2 and G3 results will differ from each other from model to model, which will inform us of the effects of different treatments of stratospheric aerosols.

The radiative forcing due to anthropogenic greenhouse gases and aerosols has already been estimated in preparing the RCP4.5 runs. Therefore, in the G3 simulations, this forcing simply needs to be balanced by aerosol forcing. Hansen *et al.* (2005) found that the radiative forcing at the tropopause due to a large tropical volcanic eruption such as Pinatubo, after allowing stratospheric temperatures to adjust, is -24τ W m⁻², where τ is the sulphate aerosol optical depth at 550 nm. In their geoengineering simulations, Jones *et al.* (2010) report an increase in sulphate aerosol optical depth of 0.05 after 3–4 years, by which time the aerosol layer has reached an equilibrium thickness. By the formula of Hansen *et al.* (2005), this should correspond to a radiative forcing of -1.2 W m⁻², which is consistent with the results found in Jones *et al.* Therefore, the amount of aerosol injected to achieve the desired radiative forcing can use the formula by Hansen *et al.* (2005) as a rough guide. However, each modeling group likely will need to fine-tune this calculation.

Experiment G4 (Figure 4), similar to experiment G3, simulates a stratospheric sulphate aerosol layer. However, instead of achieving radiative balance, G4 involves injection of stratospheric aerosols at a specific constant annual rate, turned on abruptly in the year 2020. Results from this experiment will be helpful in assessing the uncertainties that can arise in estimating the impact of geoengineering when models are used to transform emission rates into concentrations. The sudden start of the aerosol injection in 2020 is meant to approximate the kind of action that might result from society's sudden perception of a climate warming




Figure 3. Schematic of experiment G3. The experiment approximately balances the positive radiative forcing from the RCP4.5 scenario by an injection of SO₂ or sulphate aerosols into the tropical lower stratosphere.




Figure 4. Schematic of experiment G4. This experiment is based on the RCP4.5 scenario, where immediate negative radiative forcing is produced by an injection of SO₂ into the tropical lower stratosphere at a rate of 5 Tg per year.

‘emergency’ (e.g. an immediate imperative to stop ice sheet melting).

We base the proposed rate of aerosol injection of 5 Tg SO₂ per year on several considerations. Several estimates (Rasch *et al.*, 2008b; Robock *et al.*, 2008) have indicated that 3–5 Tg per year of SO₂ injected into the lower stratosphere would offset a doubling of CO₂ concentration. An injection rate of 5 Tg SO₂ per year translates into 0.0137 Tg SO₂ per day, as in Crutzen (2006), Wigley (2006), and Robock *et al.* (2008). Rasch *et al.* (2008a) suggest 1.5 Tg of sulphur (~3 Tg SO₂) per year would be sufficient, but we propose using 5 Tg SO₂ per year, to reduce the global average temperature to about 1980 values. We choose to err on the larger side of this interval to maximize the signal-to-noise ratio of the climate response to geoengineering. Additionally, according to Heckendorn *et al.* (2009), previous studies used too small of an aerosol effective radius, meaning the amounts used in prior experiments will be less effective in cooling the planet than previously thought, bolstering our argument for the larger 5 Tg SO₂ per year injection.

An ensemble of simulations will be performed for each geoengineering experiment. The suggested method of generating each ensemble and the recommended sizes of the ensembles will closely align with the protocol set by CMIP5 for the simulations without geoengineering, but if our results show the size of an ensemble is insufficient to obtain statistically significant results, additional ensemble members will be generated. In experiments G2, G3, and G4, the geoengineering will be applied for only the first 50 years, but with the runs extended an additional 20 years to examine the response to a cessation of geoengineering.

In the RCP4.5 scenario, as outlined by Moss *et al.* (2008), the total radiative forcing in 2100 reaches and subsequently stabilizes at 4.5 W m⁻² (relative to pre-industrial levels). This stabilized forcing reflects a CO₂ equivalent concentration of 650 ppm. We have selected this scenario because, as noted by

Taylor *et al.* (2008), “RCP4.5 is chosen as a ‘central’ scenario. . . [and] is chosen for the decadal prediction experiments.” Using a more optimistic scenario in which rapid mitigation is implemented would result in less robust results and is thus not likely to be as illuminating. Conversely, choosing a scenario with higher radiative forcing would reflect an irrational and unsustainable path, since if society cannot effectively mitigate greenhouse gas emissions, geoengineering would be needed on a massive scale for a long period of time, due to the long atmospheric lifetime of CO₂ (Solomon *et al.*, 2009).

Wigley (2006), Matthews and Caldeira (2007), and Robock *et al.* (2008) performed simulations in which, after a period of time, they stopped geoengineering and then evaluated the resulting rapid warming. As this response has been fairly well established, it could be argued that further investigation of the results of stopping geoengineering at this time would not be particularly interesting. However, it will likely be very easy to continue experiments G2, G3, and G4 for an additional 20 years after a 50-year geoengineering period, so we suggest that this recovery period be a part of the experiments and analyses.

The intended audience of this paper is much broader than the climate scientists who will actually perform the experiments. For that reason, we have omitted many of the details which are somewhat incidental to understanding the design and aims of the experiments. We refer the reader to a technical report (Kravitz *et al.*, 2010) which more thoroughly describes the specifications under which the simulations should be run. This technical report is also expected to serve as a working document, which will discuss problems encountered and modifications to the protocol if that should become necessary.

3. Model specifications

The models used should be the same as those used in the CMIP5 simulations. A fully coupled atmosphere

and ocean general circulation model (AOGCM) is necessary for these experiments to properly assess the dynamic responses of the climate.

Each of the models will undoubtedly treat chemistry differently. Rasch *et al.* (2008b) provide a thorough discussion of the chemistry of stratospheric injection of sulphate aerosols. If a model cannot handle photochemical conversion of sulphate aerosol precursors into sulphate aerosols, an aerosol size distribution should be used. Some models will be able to inject SO₂ and calculate the resulting aerosols. Others will require the injection of aerosols, which they will then transport, or the complete specification of the aerosol distribution and radiative properties, which will be provided by the GeoMIP project team. In the G3 experiment, some models will specify a time-invariant sulphate aerosol size distribution while others (Heckendorn *et al.*, 2009) will allow the aerosols to grow over time, which, in a realistic model, will lead to a reduction of the aerosol backscattering efficiency and an increase in the sedimentation rate. The differences between model results arising from different aerosol treatments will need to be evaluated and understood.

Models with detailed treatment of stratospheric chemistry (Morgenstern *et al.*, 2010) typically are run with a specified seasonal cycle of the ocean. Their participation in GeoMIP, particularly G3 and G4, will provide us with additional evaluation of the effects of stratospheric geoengineering on ozone chemistry.

The specifications outlined here may exclude certain modeling groups from participating in this complete suite of proposed experiments. However, we encourage these groups to participate in whatever capacity they are capable and to adhere to the experiment protocol as closely as possible. Although their results may not be directly comparable, they will undoubtedly be interesting and useful to this study.

Acknowledgements

We thank Bjorn Stevens, Drew Shindell, Jerry Meehl, Ron Stouffer, Andy Jones, Jim Haywood, Phil Rasch, and Marco Giorgetta for their suggestions in improving this document and the outlined scenarios therein. We also thank the reviewers for their thorough, helpful comments. This document also benefited from extensive discussion with attendees of the Strategic Workshop on Geoengineering Research, Hamburg, Germany, 25–26 November 2009, and subsequent discussions with researchers from the IMPLICC project. We thank Luke Oman and Allison Marquardt for their past work on and assistance with our research. The work of B. Kravitz, A. Robock, and G. Stenchikov is supported by NSF grant ATM-0730452. The work of O. Boucher is supported by DECC/Defra Integrated Climate Programme (GA01101). The work of H. Schmidt and M. Schulz is supported by the European Commission within the FP7 project IMPLICC. K. E. Taylor's contribution was supported by the Department of Energy's (DOE's) Global and Regional Climate Modeling Program, and this work was performed under the auspices of the DOE at Lawrence Livermore National Laboratory under contract DE-AC52-07NA27344.

References

- Bala G, Duffy PB, Taylor KE. 2008. Impact of geoengineering schemes on the global hydrological cycle. *Proceedings of the National Academy of Sciences* **105**: 7664–7669, DOI: 10.1073/pnas.0711648105.
- Crutzen PJ. 2006. Albedo enhancement by stratospheric sulfur injections: a contribution to resolve a policy dilemma? *Climatic Change* **77**: 211–220, DOI: 10.1007/s10584-006-9101-y.
- Govindasamy B, Caldeira K. 2000. Geoengineering Earth's radiation balance to mitigate CO₂-induced climate change. *Geophysical Research Letters* **27**: 2141–2144, DOI: 10.1029/1999GL006086.
- Govindasamy B, Caldeira K, Duffy PB. 2003. Geoengineering Earth's radiation balance to mitigate climate change from a quadrupling of CO₂. *Global and Planetary Change* **37**: 157–168, DOI: 10.1016/S0921-8181(02)00195-9.
- Govindasamy B, Thompson S, Duffy PB, Caldeira K, Delire C. 2002. Impact of geoengineering schemes on the terrestrial biosphere. *Geophysical Research Letters* **29**: 2061, DOI: 10.1029/2002GL015911.
- Hansen J, Sato Mki, Ruedy R, Nazarenko L, Lacis A, Schmidt GA, Russell G, Aleinov I, Bauer M, Bauer S, Bell N, Cairns B, Canuto V, Chandler M, Cheng Y, Del Genio A, Faluvegi G, Fleming E, Friend A, Hall T, Jackman C, Kelley M, Kiang NY, Koch D, Lean J, Lerner J, Lo K, Menon S, Miller RL, Minnis P, Novakov R, Oinas V, Perlwitz Ja, Perlwitz Ju, Rind D, Romanou A, Shindell D, Stone P, Sun S, Tausnev N, Thresher D, Wielicki B, Wong T, Yao M, Zhang S. 2005. Efficacy of climate forcings. *Journal of Geophysical Research* **110**: D18104, DOI: 10.1029/2005JD005776.
- Heckendorn P, Weisenstein D, Fueglistaler S, Luo BP, Rozanov E, Schraner M, Thomason LW, Peter T. 2009. The impact of geoengineering aerosols on stratospheric temperature and ozone. *Environmental Research Letters* **4**: 045108, DOI: 10.1088/1748-9326/4/4/045108.
- IPCC. 2007a. Contribution of Working Group I to the fourth assessment report of the intergovernmental panel on climate change. In *Climate Change 2007: The Physical Science Basis*, Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds). Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA; 996.
- IPCC. 2007b. In *Climate Change 2007: Working Group III: Mitigation of Climate Change*, Metz B, Davidson OR, Bosch PR, Dave R, Meyer LA (eds). Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA; 851.
- Jones A, Haywood J, Boucher O, Kravitz B, Robock A. 2010. Geoengineering by stratospheric SO₂ injection: results from the Met Office HadGEM2 climate model and comparison with the Goddard Institute for Space Studies ModelE. *Atmospheric Chemistry and Physics* **10**: 5999–6006.
- Kravitz B, Robock A, Boucher O, Schmidt H, Taylor K, Stenchikov G, Schulz M. 2010. Specifications for GeoMIP experiments G1 through G4. Available online at http://climate.envsci.rutgers.edu/GeoMIP/docs/specificationsG1_G4.pdf [accessed November 2010].
- Lenton TM, Vaughan NE. 2009. The radiative forcing potential of different climate geoengineering options. *Atmospheric Chemistry and Physics* **9**: 5539–5561.
- Matthews HD, Caldeira K. 2007. Transient climate – carbon simulations of planetary geoengineering. *Proceedings of the National Academy of Sciences* **104**: 9949–9954, DOI: 10.1073/pnas.0700419104.
- Morgenstern O, Giorgetta MA, Shibata K, Eyring V, Waugh DW, Shepherd TG, Akiyoshi H, Austin J, Baumgaertner AJG, Bekki S, Braesicke P, Brühl C, Chipperfield MP, Cugnet D, Dameris M, Dhomse S, Frith SM, Garny H, Gettelman A, Hardiman SC, Hegglin MI, Jöckel P, Kinnison DE, Lamarque J-F, Mancini E, Manzini E, Marchand M, Michou M, Nakamura T, Nielsen JE, Oivié D, Pitari G, Plummer DA, Rozanov E, Scinocca JF, Smale D, Teyssède H, Toohey M, Tian W, Yamashita Y. 2010. Review of the formulation of present-generation stratospheric chemistry-climate models and associated external forcings. *Journal*

- of *Geophysical Research* **115**: D00M02, DOI:10.1029/2009JD013728.
- Moss R, Babiker M, Brinkman S, Calvo E, Carter T, Edmonds J, Elgizouli I, Emori S, Erda L, Hibbard K, Jones R, Kainuma M, Kelleher J, Lamarque JF, Manning M, Matthews B, Meehl J, Meyer L, Mitchell J, Nakicenovic N, O'Neill B, Pichs R, Riahi K, Rose S, Runci P, Stouffer R, van Vuuren C, Weyant J, Wilbanks T, van Ypersele JP, Zurek M, Birol F, Bosch P, Boucher O, Feddema J, Garg A, Gaye A, Ibarra M, La Rovere E, Metz B, Nishioaka S, Pitcher H, Shindell D, Shukla PR, Snidvongs A, Thornton P, Vilariño V. 2008. *Towards New Scenarios for Analysis of Emissions, Climate Change, Impacts, and Response Strategies*, Intergovernmental Panel on Climate Change: Geneva; 132. Available online at <http://www.ipcc.ch/meetings/session28/doc8.pdf>.
- Oman L, Robock A, Stenchikov G, Schmidt GA, Ruedy R. 2005. Climatic response to high latitude volcanic eruptions. *Journal of Geophysical Research* **110**: D13103, DOI: 10.1029/2004JD005487.
- Oman L, Robock A, Stenchikov GL, Thordarson T. 2006. High-latitude eruptions cast shadow over the African monsoon and the flow of the Nile. *Geophysical Research Letters* **33**: L18711, DOI: 10.1029/2006GL027665.
- Rasch PJ, Crutzen PJ, Coleman DB. 2008a. Exploring the geoengineering of climate using stratospheric sulfate aerosols: the role of particle size. *Geophysical Research Letters* **35**: L02809, DOI: 10.1029/2007GL032179.
- Rasch PJ, Tilmes S, Turco RP, Robock A, Oman L, Chen C-C, Stenchikov GL, Garcia RR. 2008b. An overview of geoengineering of climate using stratospheric sulfate aerosols. *Philosophical Transactions of the Royal Society A* **366**: 4007–4037, DOI: 10.1098/rsta.2008.0131.
- Robock A, Marquardt AB, Kravitz B, Stenchikov G. 2009. The benefits, risks, and costs of stratospheric geoengineering. *Geophysical Research Letters* **36**: L19703, DOI: 10.1029/2009GL039209.
- Robock A, Oman L, Stenchikov G. 2008. Regional climate responses to geoengineering with tropical and Arctic SO₂ injections. *Journal of Geophysical Research* **113**: D16101, DOI: 10.1029/2008JD010050.
- Solomon S, Plattner G-K, Knutti R, Friedlingstein P. 2009. Irreversible climate change due to carbon dioxide emissions. *Proceedings of the National Academy of Sciences* **106**: 1704–1709, DOI: 10.1073/pnas.0812721106.
- Taylor KE, Stouffer RJ, Meehl GA. 2008. A summary of the CMIP5 experiment design. http://cmip-pcmdi.llnl.gov/cmip5/docs/Taylor_CMIP5_design.pdf.
- Trenberth KE, Dai A. 2007. Effects of Mount Pinatubo volcanic eruption on the hydrological cycle as an analog of geoengineering. *Geophysical Research Letters* **34**: L15702, DOI: 10.1029/2007GL030524.
- Wigley TML. 2006. A combined mitigation/geoengineering approach to climate stabilization. *Science* **314**: 452–454, DOI: 10.1126/science.1131728.